

POSTER PRESENTATION



P341: Resistance factors to standards and procedures quality of care pplication in the republic of guinea

MM Diallo^{1*}, S Conde², BK Diallo³, T Diallo⁴, AV Dembele⁴

From 2nd International Conference on Prevention and Infection Control (ICPIC 2013) Geneva, Switzerland. 25-28 June 2013

Introduction

The study was organized around individual and collective factors of resistance associated with the implementation of the requirements and recommendations of standards and quality assurance standards of care in health facilities selected.

Objectives

To determine the factors influencing the implementation and dissemination of quality standards and procedures of the professional practices of care in health facilities in Guinea.

Methods

This cross-sectional study using the interview technique. It ran from a questionnaire on the operational side of the organization of the quality of care in health facilities. On the pan of the sample, this study has identified nine health facilities corresponding to the three levels of the health pyramid. The structures were stratified by type of institution (CHU, CMC, CS), a draw was able to identify the structures corresponding to the sample to be studied. This is a sample having a size of 88 subjects distributed among the strata according to their size represented by the staff working in administrative services and maternity health facilities drawn.

Results

The main results are: i) A little knowledge of good practice recommendations by staff, ii) SONU represent the training approach in quality, iii) low RBP applicability of quality of care (40%), iv) The application of the organizational

¹Focal Point RIPAQS Guinea, Obstetrician Gynecologist Ignace Deen NH, Conakry, Guinea

Full list of author information is available at the end of the article



© 2013 Diallo et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

constraints RBP is on average 50%, v) weaknesses of policy favoring surveys (20%) and vi) Low financing application of standards and procedures for care (41%).

Conclusion

It is recommended to train staff on best practices and clinical laboratory revise regulations for the implementation of quality standards and procedures, develop tools for monitoring and evaluating the implementation of quality standards and procedures and develop action research to resolve service issues.

Disclosure of interest

None declared.

Author details

¹Focal Point RIPAQS Guinea, Obstetrician Gynecologist Ignace Deen NH, Conakry, Guinea. ²DNEHS / MSHP, Conakry, Guinea. ³Physician Standards and Procedures section DNEHS, Conakry, Guinea. ⁴Ignace Deen NH, Conakry, Guinea.

Published: 20 June 2013

doi:10.1186/2047-2994-2-S1-P341

Cite this article as: Diallo *et al.*: **P341: Resistance factors to standards and procedures quality of care pplication in the republic of guinea.** *Antimicrobial Resistance and Infection Control* 2013 **2**(Suppl 1):P341.